

**First Iberian record of *Drino maroccana* Mesnil, 1951 (Diptera,
Tachinidae, Exoristinae), a parasitoid of *Streblote panda* Hübner, [1820]
(Lasiocampidae) caterpillars¹**

D. Calvo * and J. M^a. Molina *

Tachinids are one of the most important parasitoid families to biological control (Eggerton & Gaston, 1990; Driesche & Bellows, 1996). Usually they are solitary endoparasitoids of larvae of insects, especially Lepidoptera. Many species have been introduced, and successfully established in biological control programs, others are important as indigenous parasitoids (Grenier, 1988). The group is difficult, and its taxonomy is complex (Peigler, 1994), so it isn't surprising that the Iberian tachinid fauna is little known. Few keys are available to

determine species; there are surely some undescribed species, as well as many others with unknown corology (Tschorasnig *et al.*, 1997). About 510 species are currently known from the Iberian Peninsula (Tschorasnig, pers. comm.), approximately 170 of them have been recorded from Andalusia (Tschorasnig, 1992; Tschorasnig *et al.*, *op.cit.*). With this note we report the presence of a tachinid species, *Drino maroccana* Mesnil, 1951, from Andalusia (Southern Spain). This is a new record for the Iberian fauna of Diptera.

Drino maroccana was described in 1951 as a subspecies of *D. latigena* Mesnil, 1944. Differences between *D. maroccana* and *D. latigena*, as stated by Mesnil (1951) in his original description, are apical scutellar bristles erect; dark grey pulvilliæ; males with an irregular row of black setulae behind the postocular row; abdominal tergite V nearly as long as abdominal tergite IV. Herting (1984) raised the taxa to specific range. The species was previously known only from the North of Morocco: Rabat —type locality— and Sidi es Ouagi, parasitizing *S. panda* Hbn. (= *Taragama repanda*) (Lasiocampidae) larvae (Mesnil, 1951). This is the only tachinid parasitoid known from this host (Tschorasnig, *in litt.*).

This record —although not very surprising because of the faunistical similarities between Andalusia and Northern Africa— extends the distribution, and confirms the host-parasitoid association for this species. The finding increases the number of *Drino* from Spain to five species (Tschorasnig, *op. cit.*; Tschorasnig *et al.*, *op. cit.*): *D. atropivora* (Robineau-Desvoidy), *D. galii* (Brauer



Fig. 1.— *Drino maroccana*, from Andalusia (S Spain).

Fig. 1.— *Drino maroccana*, de Andalucía (S España).

¹ This research was supported by D.G. Investigación y Formación Agraria y Pesquera, Junta de Andalucía, project nº PIA 1301.2. Part of PhD thesis: "Streblote panda Hübner, [1820] (Lepidoptera, Lasiocampidae), incidencia sobre frutales y plantas ornamentales de Andalucía Occidental. Bases ecológicas y sugerencias para su control".

* Departamento de Protección Vegetal. Entomología. CIFA—"Las Torres-Tomejil". Aptdo. Oficial 41200. Alcalá del Río (Sevilla, España). E-mail: cifatorr@cap.junta-andalucia.es

& Bergenstamm), *D. inconspicua* (Meigen) y *D. vicina* (Zetterstedt). From all these, only *D. atropivora* and *D. inconspicua* have been previously recorded in Andalusia (Sierra de Cazorla y Segura; Tschorsnig, *op. cit.*).

STUDIED MATERIAL: *Drino maroccana* Mesnil, 1951.

UTM: 30STG63; em VII-2000; El Copero, Seville (Seville, Spain); 10 m; 1♂; in coll. CIFA "Las Torres-Tomejil". Observations: ex *S. panda* Hübner, [1820] (Lepidoptera; Lasiocampidae) on *Tamarix gallica* (Tamaricaceae). UTM: 29SQB92; El Cebollar, Moguer (Huelva, Spain); 60 m; em V-2000; 2♀♀; em 9-VII-2000; 4♂♂, 2♀♀; em 16-VII-2000; 1♂, 1♀; (2♂♂, 2♀♀ in coll. CIFA "Las Torres-Tomejil"; 3♂♂, 3♀♀ in coll. Staatliches Museum für Naturkunde (Stuttgart, Alemania). Observations: all ex *S. panda* collected on *Vaccinium x corymbosum* (Ericaceae).

ACKNOWLEDGEMENTS

Our thanks are due to Dr. H.-P. Tschorsnig of the Staatliches Museum für Naturkunde in Stuttgart for his help with the determination of specimens, and for his comments and information on this species.

References

- DRIESCHE, R. G. VAN & BELLOWS, T. S. Jr., 1996. *Biological Control*. Chapman & Hall. New York. USA. 539 pp.
- EGGLETON, P. K. & GASTON, K. J., 1990. "Parasitoid" species and assemblages: convenient definitions or misleading compromises? *Oikos*, 59: 417-421.
- GRENIER, S., 1988. Applied biological control with tachinid flies (Diptera, Tachinidae): A review. *Anzeiger für Schädlingskunde, Pflanzenschutz, Umweltschutz*, 61: 49-56.
- HERTING, B., 1984. Catalogue of Palearctic Tachinidae (Diptera). *Stuttgarter Beiträge zur Naturkunde. Serie A, Biologie*, 369: 1-228.
- MESNIL, L. P., 1951. 64 g Larvaevoridae (Tachininae) In: Lindner, E. (Ed.). *Die Fliegen der Paläarktischen Region*, Lief 166. E. Schweizerbart. Stuttgart: 161-208.
- PEIGLER, R. S., 1994. Catalog of parasitoids of Saturniidae of the world. *Journal of Research on the Lepidoptera*, 33: 1-121.
- TSCHORSNIG, H.-P., 1992. Tachinidae (Diptera) from the Iberian Peninsula and Mallorca. *Stuttgarter Beiträge zur Naturkunde. Serie A, Biologie*, 472: 1-76.
- TSCHORSNIG, H.-P., ANDERSEN, S. & BLASCO-ZUMETA, J., 1997. New or interesting records of Tachinidae (Diptera) from the Iberian Peninsula. *Stuttgarter Beiträge zur Naturkunde. Serie A, Biologie*, 556: 1-46.

Recibido, el 5-IV-2002
Aceptado, el 17-V-2002
Publicado, el 5-IX-2002